

References

1. D'Orsi CJ, et al. Breast Imaging Reporting and Data System: ACR BI-RADS. 5th ed. Reston, Va: American College of Radiology, 2013.
2. Pisano ED, et al. Digital Mammographic Imaging Screening Trial (DMIST) Investigators Group. Diagnostic performance of digital versus film mammography for breast-cancer screening. N Engl J Med. 2005 Oct 27;353(17):1773-83.
3. Carney PA, et al. Individual and combined effects of age, breast density, and hormone replacement therapy use on the accuracy of screening mammography. Ann Intern Med. 2003 Feb 4;138(3):168-75.
4. McCormack VA, dos Santos Silva I. Breast density and parenchymal patterns as markers of breast cancer risk: a meta-analysis. Cancer Epidemiol Biomarkers Prev 2006;15(6):1159-1169.
5. Boyd NF, et al. Mammographic density and breast cancer risk: current understanding and future prospects. Breast Cancer Res. 2011; 13(6):223.
6. Chiu SY, et al. Effect of baseline breast density on breast cancer incidence, stage, mortality, and screening parameters: 25-year follow-up of a Swedish mammographic screening. Cancer Epidemiol Biomarkers Prev. 2010; 19(5):1219-1228.
7. Freer PE. Mammographic breast density: impact on breast cancer risk and implications for screening. Radiographics. 2015 Mar-Apr;35(2):302-15.
8. Sickles EA. The use of breast imaging to screen women at high risk for cancer. Radiol Clin North Am 2010;48(5):859-878.
9. Boyd NF, et al. Mammographic density and the risk and detection of breast cancer. N Engl J Med. 2007; 356(3):227-236.
10. Byrne C, et al. Mammographic features and breast cancer risk: effects with time, age, and menopause status. J Natl Cancer Inst. 1995;87(21):1622-1629.
11. Ursin G, et al. Mammographic density and breast cancer in three ethnic groups. Cancer Epidemiol Biomarkers Prev. 2003;12(4):332-338.
12. Boyd NF, et al. Quantitative classification of mammographic densities and breast cancer risk: results from the Canadian National Breast Screening Study. J Natl Cancer Inst. 1995;87(9):670-675.
13. Vacek PM, Geller BM. A prospective study of breast cancer risk using routine mammographic breast density measurements. Cancer Epidemiol Biomarkers Prev. 2004;13(5):715-722.
14. Bertrand KA, et al. Mammographic density and risk of breast cancer by age and tumor characteristics. Breast Cancer Res. 2013;15(6):R104.
15. Singletary SE. Rating the risk factors for breast cancer. Ann Surg. 2003 Apr;237(4):474-82.
16. Kelly KM, et al. Breast cancer detection using automated whole breast ultrasound and mammography in radiographically dense breasts. Eur Radiol. 2010 Mar;20(3):734-42.
17. Giuliano V, Giuliano C. Improved breast cancer detection in asymptomatic women using 3D-automated breast ultrasound in mammographically dense breasts. Clin Imaging. 2013 May-Jun;37(3):480-6.
18. Brem RF, et al. Assessing improvement in detection of breast cancer with three-dimensional automated breast US in women with dense breast tissue: the Somolnsight Study. Radiology. 2015 Mar;274(3):663-73.
19. Wilczek B, et al. Using 3D automated breast ultrasound to mammography screening in women with heterogeneously and extremely dense breasts: Report from a hospital-based, high-volume, single-center breast cancer screening program. Eur J Radiol. 2016 Sep;85(9):1554-63.

ABUS Locations

Century Park

201, 2377 111 St NW
Edmonton

Hys Centre

202, 11010 101 St NW
Edmonton

Summit Centre

102, 200 Boudreau Rd
St Albert AB

Terra Losa

9566 170 St NW
Edmonton



For some women, peace of mind means looking beyond mammography.

Learn how ABUS can help you offer better breast cancer screening to your patients with dense breast tissue.

Visit mic.ca



FAQs about Breast Density and Automated Breast Ultrasound (ABUS)

1 What is breast density?

Breast density is determined by the radiologist interpreting the mammogram and is assigned into one of four possible categories¹ (see below). Breast density as shown on mammography depends on the extent of fibrous and glandular tissue in the breasts and has no correlation with findings on palpation.

Mammographic Breast Density Classification

Density	% of Breast Composed of Fibroglandular Tissue	ACR Category	% of Population ²
Fatty	<25%	A	10%
Scattered fibroglandular	25 - 50%	B	43%
Heterogeneously dense	51- 75%	C	39%
Dense or Extremely dense	>75%	D	8%

2 How does breast density alter the ability of mammography to detect breast cancer?

Mammography alone is only 62-69% sensitive for the detection of breast cancer in women with heterogeneously dense or extremely dense breasts compared to 82-88% sensitive in women with lower breast density.

Decreasing Mammography Sensitivity with Increased Breast Density

Density	ACR Category	Sensitivity ³ (%)
Fatty	A	88%
Scattered fibroglandular	B	82%
Heterogeneously dense	C	69%
Dense or Extremely dense	D	62%



More than 3 out of 4 women diagnosed with breast cancer in Alberta are 50 or older.

3 What is the result of the decreased sensitivity of mammography in women with dense breasts?

Dense breast parenchyma can obscure small cancers (known as the masking effect)⁴. The risk of a patient presenting clinically with an interval cancer (between screening mammograms) is up to 18-fold higher comparing women at density extremes⁵. In the only series with sufficiently long-term follow-up to address mortality from breast cancer, Chiu et al.⁶ observed a 1.9-fold risk of breast cancer death (95%CI 1.26-2.91) among women with dense breasts after adjusting for other factors.

4 How is breast density related to a women's risk of developing breast cancer?

Women with dense breasts have an increased risk of developing breast cancer that is independent to all other risks factors (such as their family history). The increased amount of epithelial and stromal elements present in dense breasts means a greater risk that a cancer may arise⁷. Women with heterogeneously dense breasts may be 1.2-1.5 times more likely to develop breast cancer where as women with extremely dense breasts may be 2.1-2.3 times more likely⁸⁻¹⁴. By comparison, having a 1st degree family member with breast cancer makes a women 1.8 or 3.3 times more likely to develop breast cancer in her lifetime depending on whether the family member was post or pre menopausal (respectively) when diagnosed¹⁵.

Breast Cancer Risk Factors⁴⁻¹²

Risk Factor	Category of Risk	Comparison Category	Relative Risk (RR)
Alcohol Intake	2 drinks per day	Nondrinker	1.2
BMI	80 th percentile (age 55 or greater)	20 th percentile	1.2
Current Age	≥65 years	< 65 years	5.8
Past History of Breast Cancer	Invasive breast cancer	No history of invasive breast cancer	6.8
Family History	1 st degree relative ≥50 years with postmenopausal breast cancer	No 1 st , 2 nd -degree relative with breast cancer	1.8
	1 st degree relative with premenopausal breast cancer	No 1 st , 2 nd -degree relative with breast cancer	3.3
Breast Density	2 nd degree relative with breast cancer	No 1 st , 2 nd -degree relative with breast cancer	1.5
	Heterogeneously dense (ACR Category C)	Average breast density	1.2 - 1.5
	Extremely dense (ACR Category C)	Average breast density	2.1 - 2.3


5 How does ABUS help cancer detection in women with dense breasts?

Increased breast density makes it easier to distinguish malignant lesions from the adjacent breast parenchyma. In average risk women with dense breasts, adding automated breast ultrasound (ABUS) to routine screening mammography increases the detection of breast cancer by 36-167%.

Mammography Cancer Detection Rates (per 1,000) Alone and With ABUS

Density	N	Mammo alone	Mammo +ABUS	Rate Change	Rate Change (%)
Kelly et al. (2010) ¹⁶	4,419	3.6	7.2	3.6	+100%
Giuliano et al. (2013) ¹⁷	3,418	4.6	12.3	7.7	+167%
Brem et al. (2015) ¹⁸	15,318	5.4	7.3	1.9	+36%
Wilczek et al. (2016) ¹⁹	1,668	4.2	6.6	2.2	+52%
(Weighted Avg)	24,823	4.9	7.9	3.0	+66%

These cancers detected by ABUS tend to be small, invasive and node negative.



6 What are the risks of ABUS?

The two risks of ABUS for a patient are the increased risk of being recalled for supplemental imaging and an increased risk of ultimately undergoing a biopsy. The largest trial comparing mammography alone to mammography with supplementary ABUS (in average risk patients) showed that patients undergoing ABUS had an 89% increase of being recalled for additional imaging and a 93% increase rate of undergoing a breast biopsy¹⁸.

For patients coming to MIC Medical Imaging, this could potentially increase our current mammography recall rate from 6% to 11%. Similarly, this could potentially increase the percentage of our screening patients ultimately undergoing biopsy from about 1% to 2%.

As a stand-alone test, ABUS had a lower false negative rate compared to mammography in women with dense breasts (15% vs. 27%, respectively)¹⁶. A very small but indeterminate number of cancers will be occult to both mammography and ABUS.

7 How should physicians incorporate ABUS into routine breast cancer screening?

All studies using ABUS to supplement screening mammography performed the tests concurrently at 1 year intervals¹⁶⁻¹⁹. Therefore for patients with dense breasts wishing to undergo screening mammography at one year intervals, yearly (concurrent) supplementary ABUS is most appropriate at present.

Patients with dense breasts that elect to undergo screening mammography every two years could undergo ABUS concurrent to mammography, alternating every year with mammography or have ABUS yearly.

8 What about using ABUS in women with non-dense breasts?

In average risk women with non-dense breasts, ABUS should not be used for supplemental imaging for two main reasons. Firstly, mammography has good sensitivity for breast cancer in women with low (ACR category A/B) breast density due to the lack of masking effect³. Secondly, there is poor ultrasound contrast between lesions and the adjacent tissues in women with non-dense breasts, hindering lesion detection.

9 What about using ABUS in screening programs for women at high risk for breast cancer?

Current guidelines recommend annual screening with both MRI and mammography for women considered high risk for breast cancer irrespective of breast density (eg. women with BRCA gene mutation). ABUS can be considered as another supplemental screening tool in certain clinical settings (eg. in women where MRI is contraindicated) but is an inferior substitute in high risk women as MRI has a higher cancer detection rate.



1 in 8 women will be diagnosed with breast cancer in their lifetime



75% of women who develop breast cancer have no family history.