



Partnered with



Central Booking Ph 780.450.1500 Toll Free 1.800.355.1755



th Care card and and are available for X-ray Windermere W 201-6103 Curre Ft. Saskatche W SouthPointe 115-9332 South Date of L.M.P:	Date: Clinic Lo	Refer to F dentifica Sherwoo Synergy V 501 Bethe 109-Main	Preparation Instriction with this for defeated Park Veliness Centre	uctions on Reverse	
th Care card and and with Care card and and with Care card with Car	Date: Clinic Lo	Refer to F dentifica Sherwoo Synergy V 501 Bethe 109-Main	Preparation Instriction with this for defending the second	uctions on Reverse orm. St. Albert Grandin X-Ray (X-ray only 1 St. Anne ST © Summit Centre	
th Care card and and are available for X-ray Windermere W 201-6103 Curre Ft. Saskatche SouthPointe 115-9332 South	Clinic Lo	Refer to F dentifica Sherwoo Synergy V 501 Bethe 109-Main	Preparation Instriction with this found Park Vellness Centre I DR Clinic	uctions on Reverse Trm. St. Albert Grandin X-Ray (X-ray only 1 St. Anne ST Summit Centre	
th Care card and and are available for X-ray Windermere W 201-6103 Curre Ft. Saskatche SouthPointe 115-9332 South	Clinic Lo ther piece of i	Refer to F dentificar Sherwoo Synergy V 501 Bethe 109-Main	Preparation Instriction with this for defeated Park Vellness Centre I DR Clinic	uctions on Reverse orm. St. Albert Grandin X-Ray (X-ray only 1 St. Anne ST Summit Centre	
th Care card and and are available for X-ray Windermere W 201-6103 Curre Ft. Saskatche SouthPointe 115-9332 South	ther piece of i	dentifica Sherwoo Synergy V 501 Bethe 109-Main	tion with this fo	St. Albert Grandin X-Ray (X-ray only 1 St. Anne ST Summit Centre	
Windermere WW 201-6103 Curre Ft. Saskatche SouthPointe 115-9332 South	nts DR NW wan	Sherwoo Synergy V 501 Bethe 109-Main	d Park Vellness Centre I DR Clinic	St. Albert Grandin X-Ray (X-ray only, 1 St. Anne ST Summit Centre	
 Windermere 201-6103 Curre Ft. Saskatche SouthPointe 115-9332 South 	nts DR NW wan	Synergy V 501 Bethe 109-Main	Vellness Centre I DR Clinic	Grandin X-Ray (X-ray only 1 St. Anne ST Summit Centre	
Date of L.M.P:				Sturgeon Medical Women's Imaging 110-625 St. Albert Trail	
Date of L.M.P.			Stat Report	Instructions	
Date of L.M.P: Pregnant: O Yes O No Patient's Signature:			O STAT fax report O STAT verbal report to #:		
			O Send copy of X-rays with the patient		
### derangement; #### Approximate OR OL Sho OR OL Elbo ON OL Wri OD OR OL Wri OD OR OL Fing OR OL Fing Oppler OR OL Hin	derangement) Approximate date of Injury OROL Shoulder OROL Elbow: ODistal Biceps OOMedial O Latera OROL Wrist: ODOrsal O Volar ORAdial O Ulnar B* OROL Fingers: O Capsular Ligame		acute: OROL Knee: (MRI required for ACL/P cartilage and menisci) OROL Baker's cyst OROL Ankle: OAchilles O Medial OLateral OAnterior OROL Foot: Ganglion O Plantar Fascia O Morton's Neuroma		
des BPP)* O Anterior O L		· · · · · · · · · · · · · · · · · · ·		cation)	
Gastrointestinal In	naging	Bone	e Densitometr	ry	
& Duodenum)	S & D (Esophagus, Stomach Duodenum)		O Bone Densitometry O Thoracic and Lumbar Spine (Correlative x-rays) Pain Management Injection site: (eg. hip, facet, etc.)		
Whole Body Cor	nposition	T			
Requistion for these 3	_(eg: hip, knee) exams:	Blood Altern Requi	Thinners? nately, please refer	O Yes O No to our Pain Management	
	Musculoske (MRI is more a derangement) Approximate OROLShoo OROLEIbo OM S*OROLWris OD ORA Aning*OROLFing BYP)*OROLHips Other: Gastrointestinal Im OEsophagus OE, S&D (Esophagus, &Duodenum) OSmall bowel follow the OWhole Body Com Offor prosthesis loosening Requistion for these 3 asision Imaging with Ejectic	Musculoskeletal Ultrasou (MRI is more appropriate for get derangement) Approximate date of Injury in OROL Shoulder OROL Elbow: OBistal Biceps OF OMedial OLateral of ROL Wrist: OBORSAL OVOIAT ORADIAL OUTPOINT O	Musculoskeletal Ultrasound - May (MRI is more appropriate for general joint derangement) Approximate date of Injury if acute: OROL Shoulder OROL Elbow: OBISTAL BICEPS OTTICEPS OMEDIAL OLATERAL OPLETON OROL Wrist: ODORSAL OVOIAT ORADIAL OLATERAL OBORNAMENT OROL FINGERS: OTTIGGER FINGER OGANGLION OCAPSULAR LIGAMENTS (digit) OPLETON OROL HIP: OANTERIOR OLATERAL OISCHILL (HAMSTRINGS) OTHER: Gastrointestinal Imaging OESOPHAGUS O	Musculoskeletal Ultrasound - May include X-ray. (MRI is more appropriate for general joint assessment, non-siderangement) Approximate date of Injury if acute: OROL Shoulder OROL Kn ODistal Biceps OTriceps OMedial OLateral OROL Bal Note: OROL Wrist: OROL An ODOrsal OVolar ORADIA OUlnar OL OROL Fingers: OROL For OCAPSULAR (Igaments (digit)) OCAPSULAR (Igament	

Practitioner's Name:___ Practitioner's Address: Clinic Ph:_____ Clinic Fax: ____ Copy to: _____ Fax Copy: ____

Signature:

Practitioner's Stamp & Practice ID

Official Diagnostic Imaging Provider for:









EXAM PREPARATION



Ph 780.450.1500 Toll Free 1.800.355.1755 Fax 780.450.9551 | mic.ca

ALL EXAMINATIONS Please bring your Health Care card and another piece of identification with this form.

If you have any questions about your exam, exam preparation or need to change or cancel your appointment, please contact Central Booking. Patients who miss their appointment and fail to cancel 24 hours prior to their exam may be charged a \$25.00 fee.

Only exams below require preparation. Please review carefully to ensure the best exam results.

Ultrasound

O Abdomen, AAA Screen, Liver elastography Nothing to eat or drink after midnight.

O Pelvic, Renal, Bladder, Nuchal Translucency Screening, or Obstetrical (You may continue to eat)

- Empty your bladder.
- 2. Drink 1 litre of water.
- 3. Finish drinking the full amount one hour prior to the examination.
- 4. Do not empty your bladder again prior to the examination.
- O **Obstetric > 28 weeks includes BPP** (*Please have a snack prior to the exam*)
 - 1. Empty your bladder.
 - 2. Drink 500ml of water.
- 3. Finish drinking the full amount one hour prior to the examination.
- 4. Do not empty your bladder again prior to the examination.

O Abdomen with Pelvic or RLQ (Appendix)

- Nothing to eat after midnight.
- 2. The day of your exam, empty your bladder, then drink 1 litre of water.
- 3. Finish drinking the full amount one hour prior to the examination.
- 4. Do not empty your bladder again prior to the examination.
- O Anal Sphincter Exam is done to assess tears of the anal sphincter. Exam is performed both vaginally and transperineal. No preparation required.

Gastrointestinal Imaging

*If there is any chance of pregnancy, the exam should be postponed until the start of menses or within the 10 days thereafter.

O Esophagus, Stomach and Duodenum and/or Small Bowel

Do not eat or drink anything, including water, after midnight the night before your examination (if your exam is scheduled after 1:00 p.m., you may have 1 slice of dry toast and 1 cup of clear liquid prior to 7:30 a.m.). Small bowel - Time for examination varies, but may take as long as 4 hours.

Mammography

*If there is any chance of pregnancy, the exam should be postponed until the start of menses or within the 10 days thereafter.

Do not use perfume, deodorant, antiperspirant or talcum before the examination. If you experience premenstrual breast tenderness, you may delay booking until tenderness has subsided. Wear a two piece outfit. At the time of booking, you will be asked where your previous mammogram was done.

Nuclear Medicine

*If there is any chance of pregnancy, the exam should be postponed until the start of menses or within the 10 days thereafter.

Exam	Preparation	Approximate Exam Time
Gallium Scan	No preparation prior to injection. Involves 2 separate visits: O The first for injection O Two days later for images, which takes 1 hour.	15 minutes, return 48 - 72 hours later for 1 hour
Hepatobiliary Scan (HIDA)	Nothing to eat or drink after midnight	2 hours
Meckel's Scan	Starting at 8:00am the <u>day prior</u> to exam take oral Pepcid AC 20 mg (famotidine). A second 20 mg dose should be taken the evening before the exam, and a third 20mg dose taken on the morning of the test with a small sip of water, 1 hour prior to imaging. Total 3 doses. Nothing to eat or drink after midnight.	1 hour
Renal Scan: (Diuretic)	Drink 1 liter of fluid 1 hour prior to exam	Ranges from 45 minutes to 2 hours
Renal Scan: (Captopril)	Patient off ACE inhibitors for 48 hours. No breakfast. Drink 1 liter of fluid 1 hour prior to exam. Take 50mg of CAPTOPRIL 1 hour prior to examination as prescribed by your own practitioner.	1 hour
Cardiac Examinations	Refer to preparation instructions on Cardiac Requisition.	

Locations Hours of operation vary by examination

© Extended Hours available for X-ray

Edmonton

Allin Clinic (X-ray only) B1, 10155 120 ST NW Fax: 780.488.0238

() Century Park

201-2377 111 ST NW Fax: 780.461.8524 College Plaza

7th Flr-8215 112 ST NW Fax: 780.439.9977

Gateway Clinic

107-6925 Gateway BLVD NW

Fax: 1.866.815.1715

Hys Medical Centre 202-11010 101 ST NW

Fax: 780.424.7780 Namao 160

209-15961 97 ST NW Fax: 1.877.433.9020

(Tawa Centre

200-3017 66 ST NW Fax: 780.461.7527

Terra Losa

9566-170 ST NW

() Windermere

201-6103 Currents DR NW Fax: 1.888.442.2136

MRI & CT

(Separate requisition required)

MRI & CT Century Park

201-2377 111 ST NW Fax: 780.433.7286

MRI College Plaza 7th Flr-8215 112 ST NW Fax: 780.433.7286

Ft. Saskatchewan

SouthPointe

115-9332 Southfort DR Fax: 780.392.1269

Sherwood Park

Synergy Wellness Centre 501 Bethel Dr 109-Main Clinic 145-Women's Imaging

Fax: 780.392.1268 St. Albert

Grandin X-Ray (X-ray only) 1 St. Anne ST Fax: 780.458.9096

Sturgeon Medical Women's **Imaging**

110-625 St. Albert Trail Fax: 1.866.215.9996

() Summit Centre

102-200 Boudreau RD Fax: 780.459.2376

MIC Business Office Hvs Centre

203-11010 101 ST NW, Edmonton

Fax: 780.425.5979